at www.sasaweb.com

# SAJAA CPD ANSWER FORM – January/February 2021

### PLEASE SUBMIT ONLINE BEFORE 31 MARCH 2022

### Please answer the following questions:

- Submitral aneurysms usually occur in relation to:
- The posterior mitral annulus
- The anterior mitral annulus b.
- Not commonly associated with the anterior or posterior mitral annulus
- 2. Patients with submitral aneurysms usually present with:
- Mitral valve stenosis
- b. Mixed mitral valve disease
- Mitral regurgitation
- Mitral valve vegetations
- 3. An association between and submitral aneurysms has been reported. a.
- b. HIV, TB and rheumatic heart disease
- Collagen disease
- Infective endocarditis ld.
- Modified Simpson's method for determining stroke volume and cardiac output can be 4. inaccurate due to:
- Underestimation of left ventricular end diastolic volume
- Overestimation of left ventricular end systolic volume b.
- Underestimation of left ventricular end systolic volume
- 5.
- Regarding South African anaesthetic nurse training:
  Anaesthesia nurse training programmes are only offered in private nursing institutions
- No formal or accredited anaesthetic nurse training programme exists currently lb.
- Undergraduate curricula closely resemble international teaching programmes
- Regarding participants' undergraduate training: 6.
- The majority of participants stated that no or minimal anaesthesia modules were taught, and the theory was insufficient for their daily practice
- b. Participants trained at private nursing colleges stated that sufficient anaesthesia modules
- Participants stated that though minimal anaesthesia modules were taught in nursing college, the theory learnt was sufficient for their daily practice
- 7. The majority of study participants indicated that the anaesthetist they work with:
- Does have an impact on their daily work performance in theatre
- $\label{eq:makes} \mbox{Makes no difference to their daily work performance in the atre}$ b.
- Should preferably speak the same language as the anaesthetic nurse
- 8. If a nurse's theatre preparation was felt to be inadequate by the anaesthetist:
- The majority of anaesthetists showed the anaesthetic nurse how to improve their performance
- b. Only 20% of anaesthetists provided corrective teaching to the anaesthetic nurse
- The anaesthetist generally requested a senior anaesthetic nurse to assist with future cases Teaching modalities that were favoured by participants included: 9.
- Lectures at nursing college with mostly nursing tutors
- b. Online short courses by anaesthetists and nursing tutors
- Being taught by an anaesthetist and practical group sessions

### 10. The use of transparent intubation boxes:

- Significantly prolongs the time to successful intubation of a part-task trainer
- b. Should be used in patients at risk of difficult intubation during the COVID-19 pandemic, where time is of the essence
- Affects the best glottic view obtained during intubation of a part-task trainer
- Stops the spread of aerosol into the surrounding environment

### PLEASE COMPLETE QUESTIONNAIRE ONLINE

### SASA members please visit www.sasaweb.com and follow these steps:

- 1. Login with your SASA username and password under SASA Membership.
- 2. Click on the Continuing Professional Development (CPD) section.
- 3. Choose the required CPD questions.
- 4. Complete the questionnaire and click on submit.
- Your certificate is automatically generated. Select My CEUs to view and print.

Please note this is a FREE service only to members of SASA and subscribers of the journal.

#### 11. The United States Food and Drug administration has recommended that barrier enclosures without negative pressure:

- Should be maintained during intubation attempts despite difficulty with intubation
- Can be individualised to a specific subset of patients, with a low threshold to abandon the use of the enclosure
- Can be individualised to a specific subset of patients, regardless of the need for complicated airway instrumentation procedures
- Should be used in all COVID-19 patients requiring intubation, to protect healthcare workers during this pandemic

### 12. The addition of videolaryngoscopy when intubating a part-task trainer:

- Significantly reduced the time to successful intubation
- Improved the best glottic view obtained compared to direct laryngoscopy, whether or not an intubation box was used
- Increases the mouth-to-mouth distance when intubating and is therefore considered valuable
- Significantly altered the time to successful completion of intubation

#### The seniority of the intubator:

- Significantly altered the time to successful intubation (TTI)
- Significantly altered the best glottic view obtained
- Meant that 25% more consultant anaesthesiologists felt proficient at using videolaryngoscopy as compared to registrars
- Did not alter TTI or best glottic view obtained

#### Dexmedetomidine has been shown to be of particular use in the ICU setting with the following condition:

- Hypoactive delirium
- Agitated delirium
- Opioid dependence
- Thyrotoxicosis

#### 15. Early deep sedation in ICU is associated with:

- Shorter intubation periods
- An increased incidence of ventilator dyssynchrony
- Higher 180-day mortality
- Improved muscle density at 30 days d

#### 16. Duchenne muscular dystrophy has an incidence in the general population of:

- 1:3 500
- b. 1:10 000
- c. d. 1:100 1:350

### The perioperative considerations for Duchenne muscular dystrophy include:

- Malignant hyperthermia risk
- Anaesthesia-induced rhabdomyolysis risk
- Resistance to succinylcholine, necessitating a higher dose
- Avoidance of propofol

### 18. Rocuronium should be kept:

- In a maintained cold-chain of  $2-8\,^{\circ}$ C, and once removed may be kept unopened at sub-room temperatures for up to 60 days
- In a maintained cold-chain of 0-10 °C, and once removed may be kept unopened at sub-room temperatures for up to 120 days
- In a maintained cold-chain of 2-8 °C, and once removed may be kept unopened at sub-room temperatures for up to 120 days
- In a maintained cold-chain of 0-10 °C, and once removed may be kept unopened at sub-room temperatures for up to 60 days

### Rocuronium is a popular choice for:

- Elective sequence inductions, but not as an alternative muscle relaxant for modified rapid sequence inductions
- . Modified rapid sequence inductions but not for elective sequence inductions
- Elective sequence inductions as well as a commonly used alternative muscle relaxant for modified rapid sequence inductions

#### 20. In clinical practice, sealed unused rocuronium vials are often:

- Discarded after the theatre list is done
- h Kept in the theatre for emergency cases
- Returned to the pharmacy
- Returned to the refrigerator after a theatre list

## Subscribers and other recipients of SAJAA visit our new CPD portal at www.mpconsulting.co.za

- Register with your email address as username and MP number with seven digits as your password and then click on the icon "Journal CPD".
- · Scroll down until you get the correct journal. On the right hand side is an option "ACCESS". This will allow you to answer the questions. If you still can not access please send your Name and MP number to cpd@medpharm.co.za in order to gain access to the questions.
- Once you click on this icon, there is an option below the title of the journal: Click to read this issue online
- Once you have completed the answers, go back to the top of the page next to the registration option. There is another icon "Find my CPD certificate". (You will have to answer the two questions regarding your internship and last CPD audit once you have completed a questionnaire and want to retrieve your certificate).
- If you click on that icon it will open your certificate which you can print or save on your system.
- Please call MPC Helpdesk if you have any questions: 0861 111 335.



Client Support Center: +27121117001 Office - Switchboard: +27121117000

MDB015/277/01/2021