

SAJAA CPD ANSWER FORM - MAY/JUNE 2006

Subscribe to the SAJAA CPD programme and earn 16 CPD points per year

CPD reference number A008/085/03/2006

Continuing Professional Development Point Questionnaire

Accredited by the Wits CPD Office

Questions

T F

1.	HIV vascular disease manifests as aneurysmal and occlusive disease.		
2.	HAART may cause coronary artery disease.		
3.	HIV positive patients are not hypercoagulable.		
4.	Deaths due to anaesthesia may be as low as 0, 5 – 0, 8 per 100,000.		
5.	An intraoperative death has no psychological impact on the anaesthetist.		
6.	Re-expansion pulmonary oedema (REPE) never occurs after drainage of a pneumothorax.		
7.	The mortality of re-expansion pulmonary oedema is high.		
8.	Various mechanisms are proposed in the development of REPE including damage by free radicals.		
9.	All patients that are admitted to the PACU require oxygen administration.		
10.	Critical Incident Stress Debriefing should be available to anaesthetists after being involved in an intraoperative death.		
11.	The current study suggests that the typical South African seeking treatment for chronic pain tends to be female, forty years of age or older and has completed 12 years or less of formal education.		
12.	The levels of pain severity, affective distress and negative responses from significant others reported by patients in the current study are in line with the data reported for European and North American chronic pain populations.		
13.	The current study tends to suggest that chronic pain patients run a higher risk of becoming dependant upon, or addicted to medications/substances.		
14.	The minority of chronic pain sufferers report a general preoccupation with various aspects of their physical health.		
15.	Most chronic pain patients in the current study presented with significantly elevated levels of depression and anxiety.		
16.	Hallermann-Streiff syndrome has autosomal dominant inheritance		
17.	About 5% of children with Hallermann-Streiff syndrome have an underlying cardiac defect.		
18.	Children with Hallermann-Streiff syndrome may be difficult to intubate in view of their microstomia and micrognathia.		
19.	50% of children with Hallermann-Streiff syndrome are intellectually handicapped.		
20.	Hallermann and Streiff were buddies and described the syndrome together.		

SURNAME

INITIALS

MP NUMBER

Address: _____

Telephone: _____ **Fax:** _____

e-mail: _____ **Signature:** _____

Instructions

1. Read the journal to find all the answers
2. Answer the questions and mark the answer form by putting an "X" in the appropriate square.
3. Make a photocopy for your records in case the form is lost in the mail.
4. Tear out page and place the answer form in a envelope, seal it and send to:-
In House Publications, SAJAA CPD Points, PO Box 412748, Craighall, 2024
5. Do not register it and do not fax it, only original answer sheets are accepted.
6. To get your 20 CPD points subscribe to the SAJAA CPD programme. (See subscription form.)

